



NPS55-79-028

NAVAL POSTGRADUATE SCHOOL

Monterey, California





CHANNELS THAT COOPERATIVELY SERVICE A DATA

STREAM AND VOICE MESSAGES, II:

DIFFUSION APPROXIMATIONS

by

J. P. Lehoczky

and

D. P. Gaver


November 1979

Approved for public release; distribution unlimited.

Prepared for: Chief of Naval Research Arlington, VA 22717



3 76 625



Naval Postgraduate School Monterey, California

Rear Admiral T. F. Dedman Superintendent

Jack R. Borsting Provost

This report was prepared by:

Carnegie-Mellon University

Gaver, Professor

Department of Operations Research

Reviewed by:

Released by:

Michael G. Sovereign, Chairman

Department of Operations Research

William Tolles Dean of Research

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Date Entered)

REPORT DOCUMENTATION	PAGE	READ INSTRUCTIONS BEFORE COMPLETING FORM
NPS55-79-028	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
1. TITLE (and Sublitle) Channels that Cooperatively Serv	Diffusion	5 TYPE OF REPORT & PERIOD COVERED Technical nept, 6. PERFORMING ORG REPORT NUMBER
7. AUTHOR(s)	60812	8. CONTRACT OR GRANT NUMBER(s)
J. P./Lehoczky and D. P./Gaver	(16) ROCUL	1/ 12
9 PERFORMING ORGANIZATION NAME AND ADDRESS Naval Postgraduate School Monterey, Ca. 93940		PROGRAM EXEMENT PROJECT 745K 61152N RODO-01-10 N0001480WR0054
11. CONTROLLING OFFICE NAME AND ADDRESS Chief of Naval Research Arlington, VA 22217	546/ (I	NOV 379 13 NUMBER OF PAGES 43
14. MONITORING AGENCY NAME & ADDRESS(If different	nt from Controlling Office)	15. SECURITY CLASS. (of this report) Unclassified 15a. DECLASSIFICATION DOWNGRADING
Approved for public release; dis		
18 SUPPLEMENTARY NOTES 19. KEY WORDS (Continue on reverse side if necessary as		
Communication Data-V	oice Communication	ons
A system of channels mutually ac having preemptive priority but be to queue. Approximations to the	commodate both da	em, and data being allowed

DD 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

S N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Late Entered)

CHANNELS THAT COOPERATIVELY SERVICE

A DATA STREAM AND VOICE MESSAGES, II:

DIFFUSION APPROXIMATIONS*

J. P. Lehoczky Carneige-Mellon University Pittsburgh, PA

D. P. Gaver Naval Postgraduate School Monterey, CA

1. INTRODUCTION

In this paper, the study of the behavior of an element of a communication system carrying both data and voice traffic is continued. In a previous paper, Gaver and Lehoczky (1979), a fluid flow approximation was developed to predict the characteristics of the data queue length for such a system. The fluid flow approach applied to the case in which the data service rate, n, was large compared with the voice service rate, μ ; for example, when $\eta/\mu \sim 10^4$. In this paper, a different approach is taken: a Wiener process approximation is developed. The accuracy of this approximation depends on the "heavy traffic" assumption, that is that the overall traffic intensity should approach unity from below. One need not make any assumption concerning the individual parameters. It follows that the diffusion approximation complements the fluid flow approximation to give a more complete picture of the behavior of voice-data communication systems.

Research in part sponsored by ONR at Naval Postgraduate School N001480WP00067, and in part by NSF at Carnegie-Mellon University, ENG79 05526.

The communication system to be studied is essentially a part of the SENET network as described by Coviello and Vena (1975) or Barbacci and Oakley (1976). This network employs time-slotted frames. A certain portion of each frame is allocated to voice traffic, while any data traffic can use all remaining capacity including any left unused by voice. The voice traffic cannot use any unused data capacity and operates as a loss system.

We introduce probabilistic assumptions, conventional in many queueing studies. Voice traffic arrives according to a Poisson(λ) process, and each voice customer has an independent exponential(µ) service time. Data arrivals are governed by an independent Poisson(δ) process and exhibit independent exponential(n) service times. A total of c channels are reserved for exclusive use of data, while v channels can be used by both data and voice; however, voice pre-empts data. The assumptions imply that voice can be modelled as an M/M/v/v loss system, and the well-known "Erlang B" loss formula will give the loss rate. We focus our attention on the behavior of the data queue and seek to develop expressions for the steady-state distribution and the mean queue length. Many authors have studied such a system, including Halfin and Segal (1972), Halfin (1972), Fischer and Harris (1976), Bhat and Fischer (1976), Fischer (1977), Chang (1977), and Gaver and Lehoczky (1979). Only the latter paper gives approximations valid for the extreme but realistic case in which η/μ is large, and none give a Wiener approximation.

Rather than focus on this simple version of the problem, we wish to provide a somewhat more general analysis. The generalizations arise when one considers more complex voice service.

Two types of generalizations are rated here.

In general, voice traffic is unlike data traffic. A voice communication is actually a series of bursts or talkspurts separated by silence. It is reasonable to model a conversation as a two-state Markov chain of alternating talk and silence. Exponential holding times are appropriate for talkspurt and silence lengths, see Brady (1965). During the periods of silence, the voice channel is available for data traffic. This extra channel capacity can serve to allow for increased data utilization, or to reduce the data queue length, or both. The voice process can be described by a Markov chain for which each state provides a description of the number of voice customers requesting service and the number actually using the channel. For each voice state one can compute the number of channels available for data, and thus the data transition rates.

A second generalization allows for the addition of extra voice channels by reducing the quality. One strategy available to provide adequate voice and data capacity is to assign voice users to certain acceptable quality voice channels, say 6 or 8 KBPS. When a specified number of voice users are in the system, new voice users are assigned to less acceptable channels, say with 2 or 4 KBPS. In this way, one has increased the number of voice users that can be serviced at the expense of the quality of the voice

transmissions. This situation can also be incorporated into our structure. One again defines voice states: each state provides a description of the voice channels in use. One can specify a generator Q for the Markov process over the states and use it to compute the channel availability for data for each state. To include this and other possible generalizations, we assume that the voice process is a continuous time Markov chain with finite state space and that it subordinates the data queue length process. We will work out formulas for the special case in which the voice is an M/M/v/v queueing system.

2. Diffusion Approximation Approach

We assume that the voice process $\{V(t), t \geq 0\}$ can be described by a Markov chain with state space $\{1,2,\ldots,N\}$. We let this Markov chain have generator $Q=(q_{ij})$, an $N\times N$ matrix, and stationary distribution n. An important special case occurs when voice operates as an M/M/v/v loss system in which case

$$Q = \mu S = \mu$$

$$v = \frac{1}{\sqrt{1 - (1 + \rho_{v})}} \qquad \rho_{v} \qquad \rho_{v}$$

and then $\pi = (\pi_0, \dots, \pi_v)$ with

$$\pi_{i} = \frac{\rho_{v}^{i}/i!}{\sum_{j=0}^{\nu} \rho_{v}^{j}/j!}, \qquad \rho_{v} = \frac{\beta}{\mu}.$$

Acces	sion For	
	GRA&I	
DOC T		
	ounced	
Justi	fication	
By	ibution/	
	1040.0.17	
_Ava1	lability Co	Hos.
	Avail and/	or
Dist	special	1
A		

Each voice state i gives a number of channels available for data, say c_i . From c_i one can compute the rate, r_i , at which the data queue increases or decreases. Here $r_i = \delta - \eta c_i$. We let

For stability one requires $\sum_{i=1}^N \pi_i r_i < 0$. In the special case for which Q is given by (2.1), these conditions become

$$\sum_{i=0}^{\mathbf{v}} \frac{\rho_{\mathbf{v}}^{i}}{i!} (\delta - (\mathbf{c} + \mathbf{v} - i)\eta) < 0$$

where data has exclusive use of c channels and the voice has v channels. This condition can be written $\rho < 1$ where $\rho = (\rho_d + \rho_v(1-q))/(c+v), \ \rho_d = \delta/\eta, \ \rho_v = \lambda/\mu, \ \text{and} \ \ q \ \ \text{is the blocking probability}$

$$q = \frac{\rho_{v}^{v}/v!}{\sum_{j=0}^{v} \rho_{v}^{j}/j!}$$

We develop a Wiener process approximation under a heavy-traffic assumption that $\sum_{i=1}^{N} r_i \pi_i \neq 0$. This entials finding an appropriate infinitesimal mean and variance. Locally the data

queue length process does not resemble a Wiener process. If voice is in state i, the change in the data queue is nearly deterministic and equal to r_i dt rather than a normally distributed random variable. Nevertheless, if we consider a longer period of time, say [t, t+T] with T large, then the change in queue length will be more reasonably assumed to be normally distributed. The next section contains the outline of a proof that a suitably sealed version of the process converges to a Wiener process. This proof does not require the fluid flow approach of η/μ large, so a slightly more refined variance term is derived. The Wiener process derived will have a reflecting barrier at 0.

The voice process subordinates the data process; see Feller [], pp. for the notion of subordination. If voice is in state V(t), then the data queue is increasing at rate $r_{V(t)}$ over [0, t + dt]. The expected change over [t, t+T] is given by

$$E \int_{t}^{t+T} r_{V(s)} ds = \int_{t}^{t+T} E(r_{V(s)}) ds = \int_{t}^{t+T} \int_{i=1}^{N} r_{i} P(v(s)=i) ds.$$

For the case of T large we can assume V(t) has the stationary distribution, thus V(s) has distribution π for all s. The expected change over [t, t+T] thus becomes T $\sum_{i=1}^{N} r_i \pi_i$. Notice that the data queue is very large, so the boundary is not encountered. For a Wiener process, this expected change would be

mT again assuming that the boundary is not encountered. We thus find

$$\mathbf{m} = \sum_{i=1}^{N} \mathbf{r}_{i} \mathbf{\pi}_{i} = \mathbf{\pi}^{T} \mathbf{R} \mathbf{1}$$
 (2.3)

We follow a similar method to determine σ^2 . The variance in the change in the data queue length over [t, t+T] is given by

$$\operatorname{Var} \int_{t}^{t+T} r_{V(s)}^{ds} .$$

$$\begin{aligned} & \text{Var } \int_{0}^{T} r_{V(s)} \, ds \\ & = E \left(\int_{0}^{T} r_{V(s)} \, ds \right)^{2} - \left(E \int_{0}^{T} r_{V(s)} \, ds \right)^{2} = E \left(\int_{0}^{T} r_{V(s)} \right)^{2} - \left(\frac{\pi Rl}{2} \right)^{2} T^{2} \end{aligned}$$

The last term can be computed to be

$$\begin{split} E\left(\int_{0}^{T} r_{V(s)} ds\right)^{2} &= E \int_{0}^{T} \int_{0}^{T} r_{V(s)} r_{V(t)} ds dt \\ &= 2 \int_{0}^{T} \int_{0}^{T} \sum_{i=1}^{N} r_{i} P(V(s)=i) E(r_{V(t)} | V(s)=i) ds dt \\ &= 2 \int_{0}^{T} \int_{0}^{T} \sum_{i=1}^{N} r_{i} \prod_{j=1}^{N} r_{j} P(V(t)=j | V(s)=i) ds dt \end{split}$$

We let $p_{ij}(u) = P(V(u)=j|V(0)=i)$ and $P(u) = (p_{ij}(u))$. The Kolmogorov forward equations give $P(u) = \exp(Qu)$ where $\exp(M) = I + M + M^2/2! + \cdots$ for a square matrix M. One can reexpress the second moment in matrix form as

$$\begin{split} E\left(\int\limits_{0}^{T} r_{V(s)} \, \mathrm{d}s\right)^{2} &= 2 \int\limits_{0}^{T} \int\limits_{0}^{t} \underbrace{\pi_{R} \exp\left(Q\left(t-s\right)\right)}_{Q} \underbrace{Rl}_{Q} \, \mathrm{d}s \mathrm{d}t \\ &= 2 \underbrace{\pi_{R}}_{0} \int\limits_{0}^{T} \int\limits_{0}^{t} \exp\left(Qu\right) \, \mathrm{d}u \mathrm{d}t \underbrace{Rl}_{Q} \, . \end{split}$$

In view of the fact that Q is singular, one can most easily carry out the required integration by introducing the eigenvalue decomposition $Q = \stackrel{\Phi}{\Phi} D^{\Psi}$ with

$$\mathfrak{D} = \begin{pmatrix} \theta_1 & & & \\ & \theta_2 & & \\ & & \ddots & \\ & & & \theta_N \end{pmatrix}$$

a diagonal matrix of eigenvalues,

$$\Phi = (1, \Phi_1, \dots, \Phi_N), \qquad \Psi = (\pi, \Psi_2, \dots, \Psi_N)^T$$

the associated right- and left-eigenvectors. Here $\underbrace{\phi\psi}_{i} = \underbrace{\psi\phi}_{i} = I$. Since \underbrace{Q}_{i} is irreducible and finite θ_{i} has a negative real part for $2 \le i \le N$. It follows that

$$\int_{0}^{T} \int_{0}^{t} \exp(Q_{u}) dudt = \int_{0}^{T} \int_{0}^{t} \exp(Du) dudt$$

$$\psi = 2\Phi \left(\begin{array}{cccc} & & & & & \\ & \alpha_2 & & & \\ & & \ddots & \\ & & & \ddots & \\ & & & & \alpha_N \end{array} \right) \quad \stackrel{\Psi}{\sim} \quad$$

with $\alpha_i = \theta^{-i}(\theta^{-i}(e^{\theta_i^T}-1))$. Since θ_i has a negative real part $\alpha_i \sim -T/\theta_i$ as $T \to \infty$. For large T, the second moment can be written approximately as

thus the variance is given by

$$T^{2}\pi^{+}R \qquad \begin{pmatrix} 1 & & & \\ & 0 & & \\ & & & \\ & & & \\ & & & \\ & & & \\ \end{pmatrix} \underbrace{\begin{array}{c} \Psi_{R1} - 2T\pi^{T}R\Phi \\ & \Psi_{R1} - 2T\pi^{T}R\Phi \\ & &$$

The first and last terms cancel to give

$$\operatorname{Var}(\int_{0}^{T} r_{V(s)} ds) = -2T \pi^{T} \Re \Phi \begin{pmatrix} 0 & \theta_{2}^{-1} & & \\ & \ddots & & \\ & & \ddots & \\ & & & \theta_{N}^{-1} \end{pmatrix} \qquad (2.4)$$

For a Wiener process, this variance would be given by To^2 . We let

$$Q^{-} = \bigoplus_{v} \begin{pmatrix} 0 & & & & & \\ & \theta_{2}^{-1} & & & \\ & \ddots & & \\ & & \ddots & & \\ & & & \theta_{N}^{-1} \end{pmatrix} \stackrel{\Psi}{\approx}$$
 (2.5)

so Q^- is a reflexive generalized inverse of Q obtained by reciprocating nonzero eigenvalues. We note that both $QQ^-Q = Q$ and $Q^-QQ^- = Q^-$ hold, thus we write

$$\frac{\sigma^2}{2} = -\pi^T RQ^- R1$$

$$m = \pi^+ R1 < 0 .$$
(2.6)

The crucial parameter $2m/\sigma^2$ can thus be calculated.

We illustrate the formulas in (2.6) for the special case where Q is given by (2.1). For this case

$$= \frac{1}{\mu} \pi^T RS^-(R - mI) 1 + \frac{m}{\mu} \pi^T RS^-I.$$

For our choice of S, S1 = 0, thus

$$\frac{\sigma^2}{2} = \frac{1}{\mu} \pi^T \Re S^- (\Re - mI) = \frac{1}{\mu} \pi^T \Re S^- \chi$$

where y = (R - mI)1. We note that $\pi^T y = 0$, so if $z = \sqrt[3]{y}$, then z = y is a consistent set of equations and $\pi^T z = 0$.

In general one can solve (2.6) by direct calculation of Q^- ; however, this is rarely possible. A second approach is to solve Qz = y subject to $\pi^Tz = 0$. For Q of form (2.1), this can be carried out. The equations become

$$\rho_{v}(z_{i} - z_{i-1}) - (i-1)(z_{i-1} - z_{i-2}) = r_{i-1} - m, 1 \le i \le v$$

$$-v(z_{v} - z_{v-1}) = r_{n} - m$$

$$\sum_{i=0}^{v} \frac{\rho_{v}^{i}}{i!} z_{i} = 0.$$
(2.7)

One can easily find

$$z_k - z_{k-1} = \frac{\sum_{j=0}^{k-1} \pi_j (r_j - m)}{\rho_v \pi_{k-1}}$$
.

We define

$$T_{\mathbf{k}} = \sum_{j=0}^{\mathbf{k}} \pi_{j} (\mathbf{r}_{j} - \mathbf{m}).$$

Thus $T_v = 0$ and $z_k - z_{k-1} = T_{k-1}/\rho_v \pi_{k-1}$. Moreover

$$z_k = z_0 + \sum_{j=1}^k z_j = z_0 + \sum_{j=1}^k \frac{T_{j-1}}{\rho_v^{\pi}_{j-1}}$$
.

The $\sum_{i=0}^{v} \pi_i z_i = 0$ condition gives

$$z_0 = -\sum_{k=1}^{v} \pi_k \sum_{j=1}^{k} \frac{T_{j-1}}{\rho_v \pi_{j-1}}$$
.

The vector $\underset{\sim}{\mathbf{z}}$ has now been explicitly determined, and it remains to compute

$$\frac{1}{\mu} \pi^{T} R z = \frac{1}{\mu} \pi^{T} R (z_{0} + (0, u_{1}, \dots, u_{v}))$$

with

$$u_k = \sum_{j=1}^k \frac{T_{j-1}}{\rho_v T_{j-1}}.$$

We find

$$\frac{1}{\mu} \pi^{T} R z = -\frac{m}{\mu} z_{0} + \frac{1}{\mu} \sum_{k=1}^{V} \pi_{k} r_{k} \sum_{i=1}^{k} \frac{T_{i-1}}{\rho_{v} \pi_{i-1}}$$

$$= \frac{1}{\mu} \sum_{k=1}^{V} \pi_{k} (r_{k} - m) \sum_{i=1}^{k} \frac{T_{i-1}}{\rho_{v} \pi_{i-1}}$$

$$= \frac{1}{\mu} \sum_{i=1}^{V} \frac{T_{i-1}}{\rho_{v} \pi_{i-1}} \sum_{k=i}^{V} \pi_{k} (r_{k} - m) = \frac{1}{\mu} \sum_{i=1}^{V} \frac{T_{i-1}}{\rho_{v} \pi_{i-1}} (-\sum_{k=0}^{i-1} \pi_{k} (r_{k} - m))$$

$$= -\frac{1}{\mu} \sum_{i=1}^{V} \frac{T_{i-1}^{2}}{\rho_{v} \pi_{i-1}} = -\frac{1}{\mu} \sum_{i=0}^{V-1} \frac{T_{i}^{2}}{\rho_{v} \pi_{i}}.$$

We thus derive

$$\frac{\sigma^2}{2} = \frac{1}{\mu} \sum_{i=0}^{v-1} \frac{T_i^2}{v_i} , \quad T_i = \sum_{j=0}^{i} \pi_j (r_j - m)$$
 (2.8)

For the special case considered $r_j = S - (c+v-j)\eta$ = $\eta (\rho_d - (c+v) + j)$. One finds

$$m = \sum_{i=0}^{v} \eta (\rho_{d} - (c+v)+j) \frac{\rho_{v}^{j}}{j!s_{v}}$$
$$= \eta [\rho_{d} - (c+v) + \rho_{v}(1-q)]$$

where $q = \rho_V^V/v!S_V$ is the blocking probability. Thus $m = -\eta (c+v) (1-\rho)$. Plugging into (2.8) we find

$$\frac{\sigma^{2}}{2} = \frac{\eta^{2}}{\rho_{\mathbf{v}}^{\mu}} \sum_{i=0}^{\mathbf{v}-1} \frac{(\sum_{j=0}^{i} \frac{\rho_{\mathbf{v}}^{j}}{j!} [\rho_{\mathbf{d}} - (c+\mathbf{v})\rho + j])^{2}}{S_{\mathbf{v}} \rho_{\mathbf{v}}^{i} / i!}.$$

This can be simplified by letting $q_i = \rho_v^i/i!S_i$, the blocking probability for an M/M/i/i system with traffic intensity ρ_v . We find

The special case v = 1 gives

$$\frac{\sigma^2}{2} = \frac{\eta^2}{\mu} \frac{\rho_V}{(1 + \rho_V)^2} .$$

We thus have given explicit formulas for the infinitesimal mean and variance of a Wiener process with reflecting barrier at 0. One can therefore use standard results for Wiener processes with reflecting barriers to derive the stationary distribution—it will be exponential with parameter $2m/\sigma^2$. The stationary mean queue length will thus be $\sigma^2/2m$; where σ^2 and m are given by (2.6) and (2.8). For example when v=1 and c=0

$$\frac{\sigma^2}{2m} = \frac{\eta}{\mu} \frac{\rho_{\rm v}}{(1-\rho)(1+\rho_{\rm v})^3} .$$

This is developed under heavy traffic conditions, i.e., $\rho \nearrow 1$. For such conditions

$$\rho_{d} + \frac{\rho_{v}}{1 + \rho_{v}} + 1$$

or

$$\left|\rho_{\mathbf{d}} - \frac{1}{1 + \rho_{\mathbf{v}}}\right| \to 0 \quad .$$

It follows that the above expression is also

$$\frac{\eta}{\mu} \frac{\rho_{\mathbf{v}}^{\rho} d}{(1-\rho) (1+\rho_{\mathbf{v}})^2}$$

in agreement with Fischer (1977) and Gaver and Lehoczky (1979).

The characterization of the data queue as a Wiener process with reflecting barrier allows one to discuss the dynamics of the data queue. For example, suppose the data queue is at level x at time t where x is large. One might wish to study the time that elapses until the queue becomes empty. This is the duration of a busy period started at x, and corresponds to a first-passage time for a Wiener process which we denote by t_x . Moreover one might also be interested in the area beneath the sample path until it reaches the boundary of Q, since this area gives the total time waited by all customers involved in the busy period. We let this area be A(x) and seek to compute $t_x = E(T_x)$ and a(x) = E(A(x)). Straightforward backward equation or martingale arguments give

$$t_x = -x/m$$

$$a(x) = -\frac{x^2}{2m} + \frac{\sigma^2}{2m^2} x$$
(2.10)

where m is negative, and m and σ^2 are given earlier. The distribution of the first passage time can also be easily calculated; however, its Laplace transform is most convenient to find and is given by

$$E(e^{-s\tau}x) = \exp\left[\left(\frac{x}{\sigma}\right) - \left(\frac{m}{\sigma}\right) - \sqrt{\left(\frac{m}{\sigma}\right)^2 + 2s}\right]$$
 (2.11)

4. A Refined Diffusion Approximation

In this section we continue the development of a diffusion approximation for the data queue length process. The analysis is patterned after the important semigroup methods of Burman [1979]. One develops a sequence of Markov processes and studies the behavior of their generators. One shows that the generators converge and hence concludes that the associated semi-groups converge. This entails the convergence of the finite dimensional distributions. The limiting finite dimensional distributions will be those of a certain Brownian motion. The theory underlying this approach is based on theorems of Trotter and Kato, Kurtz and Burman; the reader is referred to Burman (1979) for technical details. We will illustrate the approach in the context of the special case v = 1, c = 0. The method applies to the case of general c and v but the details are not given here.

Let $\{V(t), t \ge 0\}$ be the voice process, a Markov process with state space $\{0,1\}$. Let N(t) be the data system size at time t. We study the sequence of Markov processes $\{(X_n(t), V_n(t)), t \ge 0\}_{n=1}^{\infty}$ where

$$X_n(t) = N(nt)/\sqrt{n}$$
 and $V_n(t) = V(nt)$. (3.1)

The generator of this bivariate process is easy to compute since $V_n(t)$ subordinates $X_n(t)$. Let f(x,k) be a function with domain $[0,\infty)\times\{0,1\}$ which is smooth as a function of x for each k=0,1. The generator is given by

$$A_{n}f(x,k) = \begin{cases} \delta n[f(x + 1/\sqrt{n}, k) - f(x,k)] \\ + (1-k)nn[f(x - 1/\sqrt{n}, k) - f(x,k)] + nQf(x,k) \\ & \text{for } x \ge 1/\sqrt{n} \end{cases}$$

$$\delta n[f(1/\sqrt{n}, k) - f(0,k)] + nQf(0,k)$$

$$\text{for } x = 0.$$
(3.2)

where Q is the generator of the M/M/1/1 voice process and is given by (2.1).

Next expand in a Taylor series and collect terms to rewrite (3.2) as

where $f_{\mathbf{x}}(\mathbf{x},\mathbf{k}) = \frac{\partial}{\partial \mathbf{x}} f(\mathbf{x},\mathbf{k})$ and $f_{\mathbf{x}\mathbf{x}}(\mathbf{x},\mathbf{k}) = \frac{\partial^2}{\partial \mathbf{x}} f(\mathbf{x},\mathbf{k})$.

We wish to let $n \to \infty$ and focus attention on the data queue process alone. To derive a diffusion limit for the data

queue process alone, we introduce a sequence of functions $\langle f_n(x,k) \rangle_{n=1}^{\infty} \quad \text{for which} \quad f_n(x,k) \, \Rightarrow \, f(x) \,, \text{ where } f \quad \text{is a smooth}$ function satisfying f'(0) = 0. We wish to study the limiting behavior of $A_n f_n(x,k)$. Let

$$f_n(x,k) = f(x) + \frac{1}{\sqrt{n}} g(x,k) + \frac{1}{n} h(x,k)$$

where g and h are smooth. Clearly $f_n(x,k) + f(x)$. Substitution in (3.3) gives

$$A_{n}f_{n}(x,k) = \begin{cases} nQf(x) + \sqrt{n}[Qg(x,k) + f'(x)(\delta - (1-k)\eta)] \\ + [Qh(x,k) + g_{x}(x,k)(\delta - (1-k)\eta) + \frac{1}{2}f''(x)(\delta + (1-k)\eta)] \\ \\ x \ge 1/\sqrt{n} \end{cases}$$

$$(3.4)$$

$$nQf(0) + \sqrt{n}[Qg(0,k) + f'(0)\delta] \\ + [Qh(0,k) + g_{x}(0,k)\delta + \frac{1}{2}f''(0)(\delta)] \quad x = 0.$$

We first note that Qf(x)=0 since Q operates on the voice or k component only, not on the data. This eliminates the first terms nQf(x) and nQf(0). We next examine the $Qg(x,k)+f'(x)(\delta-(1-k)\eta)$ term. This can be rewritten as

$$\begin{split} Qg(x,k) \, + \, f'(x) \, [\delta \, - \, (1-k)\eta \, - \, ((\delta-\eta) \, + \delta \, \rho_{_{\hspace{-.1em} \boldsymbol{V}}})/(1 \, + \, \rho_{_{\hspace{-.1em} \boldsymbol{V}}}) \\ \\ + \, f'(x) \, ((\delta-\eta) \, + \delta \, \rho_{_{\hspace{-.1em} \boldsymbol{V}}})/(1 \, + \, \rho_{_{\hspace{-.1em} \boldsymbol{V}}}) \\ \\ = \, Qg(x,k) \, + \, f'(x) \, \left[- \, \frac{\rho_{_{\hspace{-.1em} \boldsymbol{V}}}}{1 \, + \, \rho_{_{\hspace{-.1em} \boldsymbol{V}}}} + \, k \right] = \, f'(x) \, \, \eta \, (1-\rho) \;\; . \end{split}$$

We now select the function g(x,k) so that

$$Qg(x,k) = -f'(x) \left[-\frac{\rho_v}{1+\rho_v} + k \right] \eta .$$

The function g(x,k) must satisfy the equations

$$\lambda (g(x,1) - g(x,0)) = f'(x) \frac{\eta \rho_{V}}{1 + \rho_{V}}$$

$$-\mu (g(x,1) - g(x,0)) = -f'(x) \frac{\eta}{1 + \rho_{V}}$$
(3.5)

The above equations are consistent and redundant, thus any g(x,k) for which

$$g(x,1) - g(x,0) = \frac{f'(x)\eta}{\mu(1 + \rho_v)}$$

will suffice. We select

$$g(x,0) = \frac{1}{2} f'(x)$$

$$g(x,1) = (\frac{1}{2} + \frac{\eta}{\mu(1 + \rho_{v})}) f'(x)$$

$$g(x,k) + (\frac{1}{2} + \frac{\eta k}{\mu(1 + \rho_{v})}) f'(x) .$$
(3.6)

or

This choice of g gives

$$A_{n}f_{n}(x,k) = \begin{cases} -\sqrt{n} \, \eta (1-\rho) \, f'(x) + [Qh(x,k) + \left(\frac{1}{2} + \frac{\eta k}{\mu(1+\rho_{V})}\right) f''(x) (\delta - (1-k)\eta) \\ + \frac{1}{2} \, f''(x) (\delta + (1-k)\eta) \, \right], & x \ge 1/\sqrt{n} \\ \sqrt{n} \left(f'(0)\delta + f'(0) \frac{\eta (\rho_{V} - k(1+\rho_{V})}{\mu(1+\rho_{V})} \right) \\ + \left[Qh(0,k) + \left(\frac{1}{2} + \frac{\eta k}{\mu(1+\rho_{V})}\right) f''(0)\delta + \frac{1}{2} \, f''(0)\delta \right] \\ x = 0 \end{cases}$$

Equation (3.7) can be rewritten recalling f'(0) = 0 as

$$A_{n}f_{n}(x,k) = \begin{cases} -\sqrt{n} & \eta(1-\rho)f'(x) + \left[Qh(x,k) + \eta f''(x) \right] \frac{1}{2} + \frac{\eta k}{\mu(1+\rho_{v})} & (\rho_{d}-(1-k)) \\ + \frac{1}{2} & (\rho_{d} + (1-k)) \right] + o(1), & x \ge 1/\sqrt{n} \end{cases}$$

$$Qh(0,k) + \eta f''(0) \left[\rho_{d} \frac{1}{2} + \frac{\eta k}{\mu(1+\rho_{v})} + \frac{1}{2} \rho_{d}\right], x = 0$$

$$(3.8)$$

or

$$A_{n}f_{n}(x,k) = \begin{cases} -\sqrt{n} & \eta(1-\rho) & f'(x) + \left[Qh(x,k) + f''(x)\eta \cdot \rho_{d}\left(1 + \frac{\eta k}{\mu(1+\rho_{v})}\right)\right] + o(1) \\ & x \ge 1/\sqrt{n} \end{cases}$$

$$Qh(0,k) + f''(0) & \eta\rho_{d}\left(1 + \frac{\eta k}{\mu(1+\rho_{v})}\right) + o(1), \quad x = 0$$

We rewrite

$$Qh(x,k) + f''(x)\eta\rho_d + \left(\frac{\eta_k}{\mu(1+\rho_v)}\right)$$

as

$$\phi_{h(x,k)} + f'(x) \frac{\eta^{2} \rho_{d}}{\mu(1+\rho_{v})} \left(k - \frac{\rho_{v}}{1+\rho_{v}} \right) + f''(x) \eta \rho_{d} \left(\frac{\eta \rho_{v}}{\mu(1+\rho_{v})^{2}} + 1 \right)$$

and choose h(x,k) so that

$$Qh(x,k) = -f''(x) \frac{\eta^2 \rho_d}{\mu(1+\rho_v)} \left(k - \frac{\rho_v}{1+\rho_v}\right)$$

The function h(x,k) must satisfy

$$\lambda (h(x,1) - h(x,0)) = f''(x) \frac{\eta^2 \rho_d \rho_v}{\mu (1 + \rho_v)^2}$$

$$-\mu (h(x,1) - h(x,0)) = -f''(x) \frac{\eta^2 \rho_d}{\mu (1 + \rho_v)^2}$$
(3.1°0)

These equations are consistent and redundant so a manifold of solutions are possible. Any one will suffice. This choice of h(x,k) allows one to rewrite (3.9) as

$$A_{n}f_{n}(x,k) = \begin{cases} -\sqrt{n} \eta(1-\rho) & f'(x) + f''(x) \eta \rho_{d}\left(\frac{\eta \rho_{v}}{\mu(1+\rho_{v})^{2}} + 1\right) + o(1) \\ & x \ge 1/\sqrt{n} \end{cases}$$

$$f''(0) \eta \rho_{d}\left(\frac{\eta \rho_{v}}{\mu(1+\rho_{v})^{2}} + 1\right) + o(1)$$
(3.11)

We now let $n \to \infty$. In order for (3.11) to converge to a sensible limit, we must invoke the heavy traffic approximation, that $\rho \to 1$. Specifically we let $\rho = \rho_n = 1 - (\theta/\sqrt{n})$ for some $\theta > 0$. Equation (3.11) becomes in the limit

$$A_{n}f_{n}(x,k) = \begin{cases} -\eta\theta f'(x) + f''(x)\eta\rho_{d}\left(\frac{\eta\rho_{v}}{\mu(1+\rho_{v})^{2}} + 1\right), & x > 0 \\ f''(0)\eta\rho_{d}\left(\frac{\eta\rho_{v}}{\mu(1+\rho_{v})^{2}} + 1\right), & x = 0 \end{cases}$$

$$f''(0) = 0$$

This is the generator of a Wiener process with reflecting barrier at 0, drift of $-\eta\theta\,,$ and

$$\frac{\sigma^2}{2} = \eta \rho_d \left(\frac{\eta \rho_v}{\mu (1 + \rho_v)^2} + 1 \right) .$$

The stationary distribution for such a process is an exponential

distribution with parameter

$$\theta \bigg/ \bigg\{ \rho_{d} \bigg(\frac{\eta \rho_{v}}{\mu (1 + \rho_{v})^{2}} + 1 \bigg) \bigg\} ,$$

thus the mean queue length is given by

$$\frac{\rho_{\mathbf{d}}}{\theta} \left(\frac{\eta \rho_{\mathbf{v}}}{\mu (1 + \rho_{\mathbf{v}})^2} + 1 \right) .$$

It is informative to try to apply this heavy traffic result to a case in which ρ is near but less than 1. One might replace θ by \sqrt{n} (1- ρ). Since we have scaled by \sqrt{n} , the stationary distribution of N(t), the unscaled queue length should be approximately exponential with parameter

$$\frac{(1-\rho)}{\rho_{d}\left(\frac{\eta\rho_{v}}{\mu(1+\rho_{v})^{2}}+1\right)}$$

The mean queue length becomes

$$\frac{\rho_{d}}{1-\rho}\left(\frac{\eta \rho_{v}}{\mu(1+\rho_{v})^{2}}+1\right).$$

This result is in exact agreement with Fischer (1977) and is therefore exact. The fluid flow approximation treats η/μ as being large, hence $\left[\eta\rho_{V}/\mu(1+\rho_{V})^{2}\right]$ is assumed to dominate 1. This gives $\left(\eta\rho_{d}\rho_{V}\right)/\left[\mu(1+\rho_{V})^{2}(1-\rho)\right]$ as the fluid flow mean queue length found in Gaver and Lehoczky (1979) and in Section 2 of this paper.

The diffusion approximation thus found represents a refinement of the fluid flow diffusion approximation. Even if η/μ is large, $\rho_{_{\bf V}}$ may be small, so the 1 may be important.

Once the Wiener process infinitesimal drift and variance have been found, one can also use the dyanmics of the Wiener process to model the dynamics of the data queue process. Busy period distributions, areas, and transient behavior in general can be determined.

BIBLIOGRAPHY

- Barbacci, M. R. and Oakley, J. D. (1976). "The integration of Circuit and Packet Switching Networks Toward a SENET Implementation," 15th NBS-ACM Annual Technique Symposium.
- Bhat, U. N. and Fischer, M. J. (1976). "Multichannel Queueing Systems with Heterogeneous Classes of Arrivals," <u>Naval Research</u> Logistics Quarterly 23
- Brady, P. T. (1967). "A Statistical Analysis of On-Off Patterns in 16 Conversations." Bell S.T.J., 73-91.
- Burman, D. Y. (1979). "An Analytic Approach to Diffusion Approximations in Queueing." Ph.D. Dissertation, New York University.
- Chang, Lih-Hsing (1977). "Analysis of Integrated Voice and Data Communication Network," Ph.D Dissertation, Department of Electrical Engineering, Carnegie-Mellon University, November.
- Coviello, G. and Vena, P. A. (1975). "Integration of Circuit/Packet Switching in a SENET (Slotted Envelop NETwork) Concept," National Telecommunications Conference, New Orleans, December, pp. 42-12 to 42-17.
- Fischer, M. J. (1977a). "A Queueing Analysis of an Integrated Telecommunications System with Priorities," INFOR 15
- Fischer, M. J. (1977b). "Performance of Data Traffic in an Integrated Circuit- and Packet-Switched Multiplex Structure," DCA Technical Report.
- Fischer, M. J. and Harris, T. C. (1976). "A Model for Evaluating the Performance of an Integrated Circuit- and Packet-Switched Multiplex Structure," IEEE Trans. on Comm., Com-24 February.
- Halfin, S. (1972). "Steady-state Distribution for the Buffer Content of an M/G/l Queue with Varying Service Rate," SIAM J. Appl. Math., 356-363.
- Halfin, S. and Segal, M. (1972). "A Priority Queueing Model for a Mixture of Two Types of Customers," SIAM J. Appl. Math., 369-379.

INITIAL DISTRIBUTION LIST

	Number of Copies
Defense Documentation Center Cameron Station Alexandria, VA 22314	2
Library Code Code 0142 Naval Postgraduate School Monterey, CA 93940	2
Library Code 55 Naval Postgraduate School Monterey, Ca. 93940	1
Dean of Research Code 012A Naval Postgraduate School Monterey, Ca. 93940	1
Attn: A. Andrus, Code 55 D. Gaver, Code 55 D. Barr, Code 55 P. A. Jacobs, Code 55 P. A. W. Lewis, Code 55 P. Milch, Code 55 R. Richards, Code 55 M. G. Sovereign, Code 55 R. J. Stampfel, Code 55 R. R. Read, Code 55 J. Wozencraft, Code 74	1 25 1 1 1 1 1 1 1
Mr. Peter Badgley ONR Headquarters, Code 102B 800 N. Quincy Street Arlington, VA 22217	1
Dr. James S. Bailey, Director Geography Programs, Department of the Navy ONR Arlington, VA 93940	1
Prof. J. Lehoczky Dept. of Statistics Carnegic Mellon University Pittsburgh, PA. 15213	5

DISTRIBUTION LIST	No. of Copies
STATISTICS AND PROBABILITY FROGRAM CFFICE OF NAVAL RESEARCH CODE 436	1
AFLINGTON 22217	
CFFICE OF NAVAL RESEARCH NEW YORK AREA CHAICE 715 BROACWAY - STH FLOOR ATIN: OR. ROBER GRAFTON NEW YORK, NY 13333	1
DIRECTOR CFFICE OF NAVAL RESEARCH ERANCH OFF 536 SCUTH CLARK STREET ATTN: DEPUTY AND CHIEF SCIENTIST CHICAGO, IL 60605	1
LIERARY NAVAL OCEAN SYSTEMS CENTER SAN DIEGO CA 92152	1
NAVY LIBRARY NATIONAL SPACE TECHNOLOGY LAB ATIN: NAVY LIERARIAN BAY ST. LGUIS MS 39522	1
NAVAL ELECTRONIC SYSTEMS COMMAND NAVELEX 320 NATIONAL CENTER NO. 1 ARLINGTON VA 20360	1
DIFECTOR NAVAL REAEARCH LABORATORY ATTH: LIERARY (DHRL) CODE 2025 NASHINGTON, C.C. 20375	1
TECHNICAL INFORMATION DIVISION NAVAL RESEARCH LABORATORY HASHINGTON, D. C. 20375	1

DISTRIBUTION LIST	No. of Copies
PRCF. C. R. BAKER DEPARTMENT CH STATISTICS UNIVERSITY OF NOTRH CARCLINA CHAPEL HILL, NORTH CARGLINA 27514	1
FRCF. R. E. DECHHOFER CEFARTMENT OF OPERATIONS RESEARCH CORNELL UNIVERSITY ITHACA NEW YORK 14850	1
FRCF. N. J. BERSHAC SCHOOL OF ENGINEERING UNIVERSITY OF CALIFORNIA IRVINE CALIFORNIA 92664	1
P. J. BICKEL CEFARTMENT OF STATISTICS LNIVERSITY OF CALIFORNIA BERKELEY, CALIFORNIA 54720	
FROF. H. W. BLOCK DEPARTMENT OF MATHEMATICS UNIVERSITY OF PITTSBURGH FITTSBURGH FA	1
PROF. JOSEPH BLUM DEPT. OF MATHEMATICS, STATISTICS AND COMPLTER SCIENCE THE AMERICAN UNIVERSITY WASHINGTON	. 1
PROF. R. A. BRADLEY DEFARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAHASSEE , FLORIDA 32306	1
FROE. R. E. BARLOW OPERATIONS RESEARCH CENTER COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BERKLEY CALIFORNIA 94720	1
MR. C. N FERNETT NAVAL COASIAL SYSTEMS LACCRATORY COUL MICH. FARAMA CITY. FLERIDA 22401 30	1

No.	οſ	Copies
-----	----	--------

PRCF. L. N. PHAT COMPUTER SCIENCE / OPERATIONS RESEARCH CINTER SOLIFERN METHEDIST UNIVERSITY DALLAS TEXAS 75275	1
FRCE. W. R. ELISCHKE DEPT. CF GLANTITATIVE BUSINESS ANALYSIS UNIVERSITY CF SCUTHERN CALIFORNIA LOS ANGELES, CALIFORNIA 90007	1
CR. DERRILL J. BCRDELON NAVAL UNCEFWATER SYSTEMS CENTER COCE 21 NEWPORT PI 02840	ı
J. E. ROYER JR DEPT. OF STATISTICS SOUTHERN METHODIST UNIVERSITY DALLAS TX 75275	1
DR. J. CHANDRA U. S. ARMY RESEARCH F. G. EOX 12211 RESEARCH TRIANGLE PARK NUFTH CARCLINA 27706	1
FREF. F. CHERNOFE DEPT. CF MATHEMATICS MASS INSTITUTE OF TECHNOLOGY CAMBRIDGE: MASSACHUSETTS 02139	1
FFOF. C. CERMAN DEFARTMENT OF CIVIL ENGINEERING AND EAGINEERING MECHANICS COLUMEIA UNIVERSITY NEW YORK 10027	1
PROF. R. L. DISNEY VIRGINIA FOLYTECHNIC INSTITUTE AND STATE UNIVERSITY DEFT. OF INCUSTRIAL ENGINEERING AND OPERATIONS RESEARCH SLACKS BURG, VA 24061	1

MR. GENE F. GLEISSNER AFFLIED MATHEMATICS LABORATORY CAVID TAYLOR NAVAL SHIP RESEARCH AND DEVELOPMENT CENTER BETHESDA MD 20084	1
PROF. S. S. CLPTA DEPARTMENT OF STATISTICS PURCUE UNIVERSITY LAFAYETTE 1ND IANA 47907	1
FROF. C. L. HANSON DEPT OF MATH. SCIENCES STATE UNIVERSITY OF NEW YORK, BINGHAMTON BINGHAMTON NY 13901	1
Prof. M. J. Hinich Dept. of Economics Virginia Polytechnica Institute and State University Blacksburg, VA 24061	1
Dr. D. Depriest, ONR, Code 102B 800 N. Quincy Street Arlington, VA 22217	1
Prof. G. E. Whitehouse Dept. of Industrial Engineering Lehigh University Bethlehem, PA 18015	1
Prof. M. Zia-Hassan Dept. of Ind. & Sys. Eng. Illinois Institute of Technology Chicago, IL 60616	1
Prof. S. Zacks Statistics Dept. Virginia Polytechnic Inst. Blacksburg, VA 24061	1
Head, Math. Sci Section National Science Foundation 1800 G Street, N.W. Washington, D.C. 20550	1

	No.	of	Copies
Dr. H. Sittrop Physics Lab., TNO P.O. Box 96964 2509 JG, The Hague The Netherlands			1
CR. R. ELASHOFF BIOMATHEMATICS UNIV. CF CALIF. LCS ANGELES CALIFORNIA 90024			
PROF. GECRGE S. FISHMAN UNIV. OF NORTH CARGLINA CUR. IN OR AND SYS. ANALYSIS PHILLIFS ANNEX CHAPEL HILL, NORTH CARCLINA 20742			1
DR. R. GNANACESIKAN EELL TELEPHONE LAB HOLPOEL, N. J. 07733			1
DR. A. J. CCLEMAN CHIEF. GR DIV. 2C5.C2, ADMIN. A428 U.S. DEPT. CF CCMMERCE WASHINGTON, E.C. 20234			1
DR. H. FIGGINS 53 BONN 1. POSTFACH 589 NASSESTRASSE 2			1
HEST GERMANY			
DR. P. T. HOLMES DEPT. OF MATH. CLEMSON UNIV. CLEMSON SCUTH CAROLINA 29631			1
Dr. J. A. Hocke Bell Telephone Labs Whippany, New Jersey 07733			1
Dr. RobertHooke Box 1982 Pinehurst, No. Carolina 28374			1

IR. D. L. ICLEHART DEPT. CF C.F. STANFCRD UNIV. STANFCRD CALIFORNIA	94305	1
Dr. D. Trizna, Mail Code 5323 Naval Research Lab Washington, D.C. 20375	•	1
Dr. E. J. Wegman, ONR, Cdoe 436 Arlington, VA 22217		1
DR. H. KGEAYASHI 18 M YCFKTCHN HEIGHTS NEW YORK		1
	10598	
CR. JOHN LEHOCZKY STATISTICS CEPARTMENT CARNEGIC-MELLEN UNIVERSITY FITTS BURCH PENNSYLVANIA	15213	1
DR. A. LEMOINE 1020 GUINCA ST. FALO ALTC. CALIFORNIA		1
	94301	
DR. J. MACCUEEN UNIV. CF CALIF. LOS ANGELES CALIFORNIA		1
	90024	
FFCF. K. T. MARSHALL DEFT. CF CF NAVAL FOSTERACUATE SCHOOL MONTEREY CALIFORNIA	53 94 0	1
DR. M. MAZUMCAR MATH. DEPT. ESTINGHOUSE RES. LABS CHURCHILL BCRC FITTSBURGH PENNSYLVANIA		1
PITISULRUH PENNSYLVANIA	15235	

	No. of Copies
PROF. W. M. FIRSCH INSTITUTE OF MATHEMATICAL SCIENCES NEW YORK UNIVERSITY NEW YORK NEW YORK 10453	1
FRIF. J. E. KACANE DEFARTMENT OF STATISTICS CAFNEGIE-MELLON FITTS EURCH . PENNSYLVANIA 15213	1
DR. RICHARD LAU EIFECTOR EFFICE OF NAVAL RESEARCH ERANCH OFF 1030 EAST GREEN STREET PASADENA	1
CA 51101	
DF. A. R. LAUFER DIRECTOR CFFICE OF NAVAL RESEARCH BRANCH OFF 1030 EAST GREEN STREET PASACENA CA \$1101	i
PROF. M. LEADBETTER DEPARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL NOFTH CARULINA 27514	à
CR. J. S. LEE J. S. LEE ASSOCIATES, INC. 2001 JEFFERSON DAVIS HIGHWAY SUITE 802 ARLINGTON VA 22202	1
FRCF. L. C. LES DEFARTMENT CA STATISTICS VIRGINIA FCLYTECHNIC INSTITUTE AND STATE UNIVERSITY BLACKSBURG VA 24061	1 .
FACE. R. S. LEVENWORTH DEPT. CE INCUSTALAL AND SYSTEMS ENCINTURING UNIVERSITY OF FUCALDA GAINSVILLE FUCALDA - 22211	1

	No. of copies
FRCE G. LIBBERMAN STANFORD INIVERSITY DEFARIMENT OF OPERATIONS RESEARCH STANFORD CALIFORNIA 94305	1
DR. JAMES R. MAAR NATIONAL SECURITY AGENCY FORT MEADE, MARYLAND 20755	1
FPCF. R. W. MAESEN DEPARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA MO 65201	1
DR. N. R. MANN SCIENCE CENTER ROCKWELL INTERNATIONAL CORPORATION F.C. BOX 1089 THOUSAND CAKS, CALIFORNIA S1360	,1
CR. W. H. MARLOW PROGRAM IN LOGISTICS THE GEORGE WASHINGTON UNIVERSITY 707 2210 STREET, N. W. WASHINGTON, D. C. 20037	1
PROF. E. MASRY DEFT. APPLIED PHYSICS AND INFORMATION SERVICE UNIVERSITY OF CALIFORNIA LA JOLLA CALIFORNIA 92093	1 .
CF. BRUCE J. MCCONALD SCIENTIFIC DIRECTOR SCIENTIFIC LIAISON GROUP OFFICE CF NAVAL RESEARCH AMERICAN EPPASSY - TUKYC AFC SAN FRANCISCO 96503	1

		No. of	copies
Dr. Leon F. McGinnis School of Ind. And Sys. Eng. Georgia Inst. of Tech. Atlanta, GA 30332			1
CR. D. R. MCNEIL DEFT. CF STATISTICS PRINCETON UNIV. FRINCETON NEW JERSEY	08540		1
DR. F. MOSTELLER STAT. CEPT. FARVARD UNIV. CAMBRICCE	00510		1
DR. M. REISER IEM IHOMAS J. WATSON RES. CTR.	02139		1
YERKTOWN HEIGHTS NEW YORK	1 0598		1
DEPT OF MATHEMATICS FOCKEFELLER UNIV. NEW YORK NEW YORK	100 21		_
DR. LINUS SCHRAGE LNIV. OF CHICAGE GRAD. SCHOOL OF BLS. 5836 GREENWOOL AVE. CHICAGC, ILLINGIS			1
Dr. Paul Schweitzer University of Rochester Rochester, N.Y. 14627	60637		1
Dr. V. Srinivasan Graduate School of Business Stanford University Stanford, CA. 94305			1
Dr. Roy Welsch M.I.T. Sloan School			1
Cambridge, MA 02139			

DISTRIBUTION LIST	No. of Copies
CR. JAKET M. MYHRE THE INSTITUTE OF DECISION SCIENCE FOR BUSINESS AND PUBLIC POLICY CLARERUIT MEN'S COLLEGE	1
CLAREMONT CA \$1711	
MR. F. NISSELSCN BLREAU OF THE CENSUS ROOM 2025 FFCERAL EVILCING 3 WASHINGTON . D. C. 2033	1
MISS B. S. CRLEANS NAVAL SEX SYSTEMS COMMAND (SEA OBF) FM 10509 ARLINGTON VIRGINIA 20369	1
FRCF. D. E DWEN DEPARTHENT OF STATISTICS SOUTHERN METHODIST UNIVERSITY CALLAS TEXAS 75222	1
Prof. E. Parzen Statistical Sceince Division Texas A & M University College Station TX 77843	1
DR. A. PETRASOVITS RCCM 2078 , FCCC AND DRUG BLDG. TUNNEY'S PASTURE CTTOWA , CNTARIC KLA-CL2 , CANADA	1
FRCF. S. L. PHCENIX SIELEY SCHOOL OF MECHANICAL AND AEROSPACE ENCINEERING CORNELL UNIVERSITY ITHACA NY 14850	1
DR. A. L. POWELL DIRECTOR OFFICE OF NAVAL RESEARCH BRANCH OFF 455 SUMMER STREET	1
BCSTCN MA 02210	
PR. F. R. FRICFI CODE 224 CRESATIONSL TEST AND UNRS EVALUATION FORCE (UPTEVEUR) NOGLILLE VINGINIA	1
20360	

DISTRIBUTION LIST	No. of Copies
PROF. H. L. PURI DEFT. EF MATHEMATICS P.G. BOX F INCIANA UNIVERSITY FOUNDATION ELCOMINGION IN 47401	1
FROF. H RCPEINS DEFARTMENT OF MATHEMATICS CCLUPETA UNIVERSITY NEW YORK: NEW YORK:	1
PFOF. M ROSENBLATT DEPARTHENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA SAN DIEGO LA JOLLA CALIFORNIA 92093	1
PROF. S. W. RCSS COLLEGE OF ENGINEERING UNIVERSITY OF CALIFORNIA BERKELEY CA 94723	1
PROF. I RUBIN SCHOOL OF ENGINEERING AND APPLIED SCIENCE UNIVERSITY OF CALIFORNIA LOS ANGELES . CALIFORNIA DO024	1
FRCF. I. R. SAVAGE DEPARTMENT OF STATISTICS YALE UNIVERSITY NEW HAVEN. CONNECTIOUT C6520	1
FRCF. L. L. SCHARF JR DEPARTMENT OF ELEGRICAL ENGINEERING COLORACU STATE UNIVERSITY FT. COLLINS, COLORACU E0521	1
PROF. R. SERFLING DEPARTMENT OF STATISTICS FLORIDA STATE UNIVERSITY TALLAMASSEE FLORIDA 32306	1
PROF. W. R. SCHUCAMY DETARTHENT OF STATISTICS SOUTHERN METHICUIST UNIVERSITY DALLAS , TEXAS 75222	1

DISTRIBUTION LIST	No. of Copies
PROF. C. C. SIEGMUND CEPT. OF STATISTICS STANFORD UNIVERSITY STANFORD	1
CA \$43.05	
FRCE. M. L. SEGOMAN DEPT. CE ELECTRICAL ENGINEERING POLYTECHNIC INSTITUTE OF NEW YORK BRCCKLYN, NEW YORK 11201	1
DR. A. L. SLAFKOSKY SCIENTIFIC ADVISOR COMMANDANT OF THE MARINE CORPS HASHINGTON, D. C. 20380	1
CR. C. E. SMITH DE SMATICS INC. P.C. BCX 618 STATE COLLEGE PENNSYLVANIA 16801	1
PROF. W. L. SPITH DEFARTMENT OF STATISTICS UNIVERSITY OF NORTH CARCLINA CHAPEL HILL NORTH CARCLINA 27514	1
Dr. H. J. Solomon ONR 223/231 Old Marylebone Rd London NW1 5TH, ENGLAND	1
MF. GLENN F. STAFLY NATIONAL SECURITY AGENCY FORT MEACE : MARYLAND 20755	1
Mr. J. Gallagher Naval Underwater Systems Center New London, CT	1
Dr. E. C. Monahan Dept. of Oceanography University College Galway, Ireland	1

DISTRIBUT	TION LIST	No. of Copies
DR. R. M. STARK STATISTICS AND COMPUTER SCI UNIV. OF DELAWARE NEWARK	•	1
CELAWARE	19711	
FFOF. RICHAR C VANSLYKE RES. ARALYSIS CORP. BEECHWOOD CLD TAFFER FO/D GLEN COVE. NEW YORK	11 542	1
PRCF. JOHN W. TUKEY FINE HALL FRINCTION UNIV. PRINCTION NEW JERSEY	·	1
	08540	
CR. THOMAS C. VARLEY CFFICE OF NAVAL RESEARCH CODE 434 ARLINGTON VA	02217	1
	22217	1
FRCE. G. WATSON FINE HALL FRINCETON UNIV. PRINCETON NEW JERSEY	C8540	•
PR. CAVIC A. SWICK ADVANCED PROJECTS GROUP		1
CODE 8103 NAVAL RESEARCH LAB. NASHINGTON	0375	
PR. WENDELL G. SYKES ARTHUR C. LITTLE, INC.		ı
ACCRIL PARK CAMBRIDGE MA 02	2140	
PROF. J. R. THOMPSON DEPARTMENT OF MATHEMATICAL S RICE UNIVERSITY HOUSTON, TEXAS 77001	SCIENCE	1
PROF. W. A. THENESCH DEFARTMENT OF STATISTICS UNIVERSITY OF MISSOURI COLUMBIA .		1
MISSOURI 65201	41	

	No. of Copies
FRCF. F. A. TILLMAN DEPT. CF INDUSTRIAL ENGINEERING KANSAS STATE UNIVERSITY MANHATTAN	. 1
€6506	
PROF J. A. TUKEY - CEFARIMENT OF STATISTICS - FRINCETON UNIVERSITY	1.
FRINCETON , N. J. 08540	
PROF. A . F . VEINOTT DEFARTMENT OF CPERATIONS RESEARCH STANFORD UNIVERSITY STANFORD CALIFORNIA 94305	į
CANIEL H. WAGNER STATION SULARE ONE FACLI, FENNSYLVANIA 19301	1
PRCF. GRACE WAHBA CEFT. CF STATISTICS UNIVERSITY CF WISCONSIN MADISON WI	1
FRCF. K. T. WALLENIUS DE FARTMENT OF MATHEMATICAL SCIENCES. CLEMSON. UNIVERSITY CLEMSON. SOUTH CARCLINA 29631	1
PROF. BERNARD WIDROW STANFORD ELECTROWICS LAB STANFORD UNIVERSITY STANFORD	1
94305	•

DISTRIBUTION LIST	No. of Copies
OFFICE OF NAVAL RESEARCH SAR IPANCISCO AREA OFFICE 760 MARKET STREET SAN FRANCISCO CALIFORNIA 94102	.1
TECHNICAL LIBRARY NAVAL CREMANCE STATION INCIAN HEAC MARYLAND 20640	1
NAVAL SHIP ENGINEERING CENTER PHILADELPHIA DIVISION TECHNICAL LIBRARY PHILADELPHIA PENNSYLVANIA 19112	1
BLREAU OF NAVAL FRESONNEL DEFARTMENT OF THE NAVY TECHNICAL LIERARY WASHINGTON C. C. 20370	1
PRCF. P. AECE'-HAMEZO DEPARTMENT OF MATHEMATICS UNIVERSITY OF NORTH CARCLINA CHARLOTTE NC 28223	1
PROF. T. W. ANCERSON DEFARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORD, CALIFORNIA '84305	1
PROF. F. J. ANSCOMBE DEPARTMENT OF STATISTICS VALE UNIVERSITY NEW HAVEN CONNECTICUT C6520	1
ROF. L. A. ARCIAN NSITIUTE OF INCUSTRIAL OFINISTRATION NION COLLUCE CLEMPCIADY . EN YORK 12308	1